

SZABADOS, Antal, dr.

"Food virology as a new field for the veterinarian-food hygienist"
by G. Seidel. Reviewed by Antal Szabados. Magy allatorv lap 17 no.8:
319 Ag '62.

I 2047 66 EWT(1)/EPE(n)-2/EWD(t) TIP(c) ID/SM/JG
ACC NR: AP6009278

SOURCE CODE: HU/0014/66/000/003/0056/0061

AUTHOR: Szabados, B. (Graduate metallurgist)

ORG: none

TITLE: Determining the gas content of molten aluminum alloys

SOURCE: Kohaszati lapok, no. 3, 1966, 56-61

TOPIC TAGS: aluminum alloy, molten alloy, alloy gas content, gas content determination, alloy fluidity, alloy shrinkage, alloy castability

ABSTRACT: The possibility of determining the gas content of liquid aluminum alloys has been tested with Dardell-type equipment in three types of alloys: Al-Si12 (12.2% Si, 0.29% Fe, 0.26% Mn); Al-Si9 (8.4% Si, 0.47% Fe, 0.36% Mg, 0.36% Mn) and Al-Mg-Si (3.05% Si, 0.24% Fe, 0.56% Mg, 0.56% Mn). The gas content of the molten alloys was measured before and after treatment with a mixture of 85% NaCl and 15% Na_3AlF_6 . The effect of the gas content on fluidity and shrinkage was evaluated. In Al-Si12 alloy at temperatures of 650 to 700°C, the gas content before salt treatment was $0.247 \text{ cm}^3/100\text{g}$ and after the treatment, $0.244 \text{ cm}^3/100\text{g}$, still above the permissible limit. Its fluidity improved only by 6.5% after salt treatment. The high value, but did not impair the strength value. In Al-Si9-Mg alloy, the salt treatment increased the fluidity by 15.6%. The gas content in the Al-Mg-Si alloy before salt

Card 1/2

DK 669.785:669.715-404

L 20434-66
ACC NR: AP6009278

treatment was too high to measure with the equipment, the tensile strength was at the lower limit, and the ductility was adequate. The low strength values were attributed to a higher iron content. The test indicated that strength characteristics are influenced by oxides rather than by gas content. Orig. art. has: 9 figures and 7 tables.

[KS]

SUB CODE: 11 / SUBM DATE: none/ ATD PRESS: 4222

Card 2/2 ULR

SZABADOS, C.; TARUS, V.

Reducing the operating costs in automotive transportation by modernizing
the methods of maintenance. I. p. 425.

REVISTA TRANSPORTURILOR. (Asociatia Stiinifica a Inginerilor si Tehnicienilor
din Romania si Ministerul Transporturilor Rutier, Navale si Aeriene)
Bucuresti, Romania. Vol. 6, no. 10, Oct. 1959

Monthly List of East European Accessions (EEAI) LC Vol. 9, no. 2, Jan 1960

Uncl.

ZARIBOS, C.; TARUS, V.

Reducing operation costs in automotive transportation by modernizing the methods of maintenance. II. p. 467.

REVISTA TRANSPORTURIILOR. (Asociatia Stiintifica a Inginerilor si Technicienilor din Romania si Ministerul Transporturilor Rutiere, Navale si Aeriene) Bucuresti, Rumania. Vol. 6, no. 11, Nov. 1959.

Monthly list of East European Accessions (EEAI) LC Vol. 9, no. 2, Feb. 1960

Uncl.

SZABADOS, C.

"Technological bases of autovehicle repairing" by R. Keller.
Reviewed by C. Szabados. Rev transport 8 no. 9:422-424
S '61.

SZMUK, Imre, dr.; BACH, Imre, dr.; DANZIGER, Laszlo, dr.; FEKETE, Balazs, dr.;
FLEISCHMANN, Laszlo, dr.; JAKO, Geza, dr.; MISSURA, Tibor, dr.;
POPPER, Szuzsanna, dr.; SZABADOS, Daisy, dr.

Use of radioiodine in localization of inflamed regions (foci,
abscesses). Orv. hetil. 97 no.34:949-951 19 Aug 56.

1. A Fovarosi Peterfy Sandor u. Korhazrendelo (igazgato:
Lendvai, Jozsef, dr.) kozlemenye.
(BRAIN, abscess
exper., localization with radioiodine in dogs (Hun))
(IODINE, radioactive
in localization of exper. brain abscesses in dogs (Hun))

SABADOS, D.

F-3

YUGOSLAVIA/Microbiology - Industrial Microbiology.

Abs Jour : Ref Zhur - Biol., No 15, 1958, 67176

Author : Sabados, D.

Inst :

Title : Bioyogurt.

Orig Pub : Mlječarstvo, 1957, 7, No 7, 145-147.

Abstract : No abstract.

Card 1/1

L 1981-66

ACCESSION NR: AT5024289

centrations of DNP, a further decrease was observed in pyruvate utilization, oxygen consumption and citrate synthesis. Both the DNP inhibition of pyruvate utilization and citrate synthesis, and its activation of acetoacetate production were diminished by small amounts of fumarate. In the presence of both substrates, DNP enhanced the oxygen consumption. Thus, in addition to its inhibiting effect on pyruvate utilization, DNP increases its conversion to acetoacetate and at the same time, especially in the presence of fumarate, renders the oxidation of acetyl-CoA (formed from pyruvate) more complete in the citric acid cycle. This means that DNP shows a ketogenic effect with respect to the pyruvate metabolism of liver mitochondria. The probable mechanism of this effect is discussed. "We are indebted to Prof. V. Szekessy-Hermann for her interest throughout this study." Orig.art. has: 1 figure, 3 tables.

ASSOCIATION: Institute of Biochemistry, University Medical School, Budapest

SUBMITTED: 00

ENCL: 00

SUB CODE: LS, OC

NR REF Sov: 000

OTHER: 0011

JPRS

Card 2/2 DP

PETER, Ferenc, dr. (Budapest VIII., Rakoczi ut 27/b); RUSZNAK, Istvan, dr. (Budapest III. Korvin O.u.44); PALYI, Gyula (Budapest XII, Meredek u.43); SZABADOS, Ida (Budapest XI., Móricz Zsigmond körter 14)

Investigation of adsorption waves. I. Investigation of the reduction of anthraquinone-1-sulphonic acid at the dropping mercury electrode. (EEAI 10:4)
Acta chimica Hung 24 no.4:363-370 '60.

1. Department for Applied Chemistry, Technical University, Budapest.
(Adsorption) (Anthraquinonesulfonic acid)
(Electrodes, Dropping mercury) (Riboflavine)
(Methylene blue) (Polarograph and polarography)
(Diffusion) (Phenazinol)

PETER, Ferenc; RUSZNAK, Istvan; PALYI, Gyula; SZABADOS, Ida

Investigation of adsorption waves. I. Examination of anthraquinone-l-sulfonic-acid reduction on mercury-dropping electrodes. Magy kem folyoir 66 no.5:178-181 My '60.

1. Budapesti Muszaki Egyetem Gyakorlati Kemial Tanszeke.

PETER, Ferenc; PALYI, Gyula; SZABADOS, Ida

Investigation of adsorption waves.II.Investigation of anthra-quinone-1-5-disulphonic acid reduction on dropping mercury electrodes. Magy kem folyoir 67 no.10:428-431 O '61.

1. Textilipari Kutato Intezet (for Peter) 2. Egyesult Vegyimuvek (for Palyi) 3. Orszagos Kozegeszsegugyi Intezet (for Szabados),

PETER, Ferenc; SZABADOS, Ida; PALYI, Gyula

Investigation of adsorption phenomena occurring on dropping mercury electrode.I.Effect of leuco-anthraquinone sulfuric acid-ester derivatives on their reduction occurring in the nitro-benzol-3-sulfonic acid agent. Magy kem folyoir 68 no.3: 101-105 Mr '62.

1. Textilipari Kutato Intezet, Budapest (for Peter) 2. Orszagos Kozegazsegugyi Intezet, Budapest (for Szabados) 3. Egyesult Vegviumuvek, Budapest(for Palyi)

PETER, Ferenc; SZABADOS, Ida

Examination of nitrobenzene-3-sulfonic acid reduction on
dropping mercury electrodes. Magy kem folyoir 68 no.4:145-149
Ap '62

1. Textilipari Kutato Intezet, Budapest (for Peter).
2. Orszagos Kozegeszseguzyi Intezet, Budapest(for Szabados).

PETER, Ferenc; PALYI, Gyula; SZABADOS, Ida

Investigation of adsorption waves.III.Reduction of anthraquinone-1,8-disulfonic acid on dropping mercury electrodes. Magy kem folyoir 62 no.6:234-236 Je '62.

1. Textilipari Kutato Intezet, Budapest (for Peter).
2. Egyesult Vegyimuvek, Budapest (for Palyi).
3. Orszagos Kozegeszsegugyi Intezet, Budapest (for Szabados).

SZABADOS, J.

POZSONYI, J.;SZABADOS, J.

Renal tuberculosis in children. Orv. hetil. 94 no. 42:1149-1152 18 Oct
1953. (CLML 25:5)

1. Doctors. 2. Osteology Department of Szabadsaghegy State Children's Sanatorium (Director - Head Physician -- Dr. Istvan Fleisch) and Urology Department of the National Physical Training and Sports Hygiene Institute (Director - Head Physician -- Dr. Sandor Balassa).

SZABADOS, Jeno, dr.

New data on antituberculotic therapy of renal tuberculosis. Magy.
sebeszet 9 no.3:191-197 June 56.

1. Az Orszagos Testneveles es Sportegeszsegugyi Intezet (igazgato:
Dr. Kovari Aladar es Fodor Jozsef Tbc. Gyogyintezet (Igazgato:
Dr. Risko. Tibor) urologiai osztalyainak kozlemenye.

(TUBERCULOSIS, RENAL, ther.

conservative ther., indic. & compl. (Hun))

SORU, Eugenia; SZABADOS, Judith; TANASESCO, D.; PARASCHIVESCO, Maria

Oxidative phosphorylation in *Mycobacterium tuberculosis* BCG and its variants resistant to streptomycin, PAS and isoniazid. Arch. roum. path. exp. microbiol. 21 no.1:59-68 Mr '62.

1. Travail de l'Institut "Dr. I. Cantacuzino" --Services de Biochimie et d'Immunochimie et du Vaccin B.C.G.
(*MYCOBACTERIUM BOVIS*) (DRUG RESISTANCE, MICROBIAL)
(ANTITUBERCULAR AGENTS) (ENERGY METABOLISM)

SZABADOS, Judith; TOPCIU, Aurica [deceased]

Research on directed resistance in *Neisseria meningitidis*. Pt.5.
Arch. Roum. path. exp. microbiol. 23 no.4:973-976 D '64.

l. Travail de l'Institut "Dr. I. Cantacuzino", Service de Bio-
chimie Generale (for Szabados), Service des Cocci Pathogenes
(for Topciu).

Szabados, Karolyne

HUNGARY/ Analytical Chemistry - Analysis of
Inorganic Substances

G-2

Abs Jour : Referat Zhur - Khimiya, No 4, 1957, 12067
Author : Bozsai Imre, Szabados Karolyne
Title : Determination of Lead Content in Lead Alloys
Orig Pub : Kohasz. lapok, 1955, 10, No 9, 423-424

Abstract : The known methods for determining Pb in alloys have been checked. The most extensively utilized molybdate method shows considerable discrepancies in the results; complete precipitation of $PbMoO_4$ takes place only at pH 8. Accuracy of Pb determination depends upon conditions of precipitation of PbS; at pH 2 no precipitation of PbS takes place, at pH 2.5 it is contaminated by iron. Optimal pH value of the medium is 2-2.5. On the basis of this verification an accurate method has been developed for the determination of Pb. Prior to treatment with hydrogen

Card 1/2

Card 2/2

SEAWAY, L.

An informative summary of the material of the Second International Conference
on the Peaceful Uses of Atomic Energy. p.25

ENERGIA ES ATOMTECHNIKA. (Energiagazdalkodasi Tudomanyos Egyesulet)
Budapest, Hungary
Vol. 12, no.1, Jan. 1959

Monthly List of East European Accessions (EEAI) IC., Vol. 8, no.7, July 1959
Incl.

SZABADOS, Laszlo, okleveles gépeszmérnök

Nuclear reactor systems. Ipari energia 1 no.1/4:63-94 J1-0
'60.

1. Kézponti Fizikai Kutató Intézet.

83323

H/008/60/000/009/002/002
B009/B05711.3000

AUTHORS:

Szentgyörgyi, István, Török, Antal, Szabados, László

TITLE:

Examination of Heat Transfer of Suspensions

PERIODICAL:

Energia és Atomtechnika, 1960, No. 9, pp. 388-395

TEXT: Some organic compounds or their mixtures (diphenyl, diphenyl, terphenyl, etc.) exhibit properties that qualify them for use as reactor coolants. Their heat-transfer capacity, however, is considerably lower than that of the conventional coolant, water. To improve the heat-transfer coefficient of these liquids, the authors suspended in them solids of high heat-transfer capacity and attained an improvement of 70% by an addition of 50 per cent by weight. The heat-transfer coefficient was calculated from Nusselt's empirical function. Heat transfer, however, also depends on viscosity. Since the viscosity of the suspension increases with respect to that of the pure liquid, the effect of the higher heat-transfer coefficient is balanced, and above a certain concentration the heat transfer of the suspension even decreases. In the graphite-diphenyl test suspension this heat transfer maximum appeared at 90°C and with a graphite addition

Card 1/2

83323

**Examination of Heat Transfer of Suspensions E/008/60/000/009/002/002
E009/B057**

of about 23 wt%. The relative viscosity versus concentration is plotted in Fig. 2 according to both Hatschek's and Orr's and Dalla Valle's formulas (Refs. 6 and 11, respectively). The experimental setup was essentially a single-tube heat exchanger. American researchers made similar experiments on graphite and aluminum suspensions (Ref. 11). For the evaluation of the test results, constants suggested by Bayer-Leverkusen, for the same mixture of diphenyl-oxide and diphenyl were substituted in the formula of the heat-transfer coefficient. The use of suspensions as reactor coolants is associated with the following disadvantages: higher power consumption of the circulating pump, higher wear and tear of pumps, metal parts, etc. The use of suspensions as reactor coolants would not be advisable even if the heat transfer could be multiplied thereby. There are 10 figures, 2 tables, and 14 references: 3 Soviet, 2 US, 1 German, 1 French, and 3 Hungarian.

ASSOCIATION: Központi Fizikai Kutató Intézet
(Central Research Institute of Physics)

Card 2/2

SZENTGYORGYI, Istvan; TOROK, Antal; SZABADOS, Laszlo

Investigations of the heat transfer properties of suspensions. Koz
fiz kozl MTA 8 no.2/3:115-129 '60. (EEAI 10:4)

1. A Magyar Tudomanyos Akademia Kozponti Fizikai Kutato Intezete,
Reaktorfizikai es Technikai Laboratorium
(Suspensions) (Nuclear reactors)
(Biphenyl) (Terphenyl)

S/262/62/000/007/001/016
I007/I207

AUTHOR: Németh, Géza, Raszl, Károly, Szabados László, Szeghő, Laszló and Torok, Antal

TITLE: Stable temperature distribution (in case of convective heat transfer) in a cylindrical fuel cell of the active zone of a heterogeneous nuclear reactor

PERIODICAL: Referativnyy zhurnal, otdel'nyy vypusk. 42. Silovyye ustavovki, no. 7, 1962, 4, abstract 42.7.13. "Magyar tud. akad. Közp. fiz. kutató int. közl.", v. 9, no. 1-2, 1961, 3-23, III, IX [Abstracter's note: Original language Hungarian].

TEXT: A solution is presented of the differential equation for convective heat transfer in finite and infinite fuel elements. For exact solutions the coolant temperature can not be reproduced without knowing the temperature distribution in each fuel element. For the solution of the given equation the temperature distribution along the fuel element is assumed to be unknown. A comparison is given between exact and approximate solutions. There are 9 figures and 13 references.

[Abstracter's note: Complete translation.]

Card 1/1

Magyar Tudományos Akadémia, Központi Fizikai
Kutató Intézete, Reaktorfizika Tanszék

26.2231

32721
H/008/62/000/001/002/002
B122/B102

AUTHORS: Nemet, G., Raszl, K., Szabados, L., Szeghő, L., Török, A.

TITLE: Steady-state heat distribution in a cylindrical-symmetric unit cell of the active zone of heterogeneous reactors in the case of convective heat transfer

PERIODICAL: Energia és Atomtechnika, no. 1, 1962, 41 - 46

TEXT: Part II. Determination of A_n coefficients. In Part I it was found that the solution of differential equation $T(r,z)$ was given by the sum of Laplace's equation (in the form of an infinite series) and of Poisson's equation (in closed form): $T(r,z) = T_0(r,z) + T_1(r,z)$ (51). In the solution of this differential equation the unknown coefficients A_n ($n = 0, 1, 2, 3 \dots$) arise. A practicable way of calculating these coefficients is the application of equation systems with an infinite number of unknowns. (Reference is made here and in the following to L. V. Kantorovich and V. I. Krylov: Approximate methods of higher analysis (Hungarian edition, Budapest, 1955)). By this method the following two Card 1/4 X

32721

Steady-state heat distribution...

H/008/62/000/001/002/002
B122/B102

equations are found to yield the system of equations with an infinite number of unknowns for the determination of the A_n :

$$A_0 = B_0 - \sum_{n=1}^{\infty} A_n \frac{r_n}{1B_n} [1 - (-1)^n] \quad (n = 1, 2, 3, \dots) \quad (73) \text{ and}$$

$$A_k = \frac{B_k}{p_k} - \frac{1}{p_k} \sum_{n=1}^{\infty} A_n \frac{2r_n \beta_n}{1(\beta_n^2 - \beta_k^2)} [1 - (-1)^{n+k}] \quad (74) \quad (n = 1, 2, \dots, (k-1), (k+1), \dots)$$

Coefficients B_0 and B_k are computable Fourier coefficients of $f(z)$. By substitution of $A_k p_k = X_k$ into equation (74), this equation is transformed to a system of entirely regular equations having (according to a thesis of Kantorovich-Krylov) but one solution which can be determined by the method of successive approximations. $B_k = O(1/k^2)$; from a thesis of the above Soviet authors it follows that $X_k = O(1/k^2)$, for the unique solution of the entirely regular equation system tends toward zero with

Card 2/4

X

32721
H/008/62/000/001/002/002
B122/B102

Steady-state heat distribution...

$k \rightarrow \infty$. In zeroth approximation $A_k^{(0)} = 0$, in first $A_k^{(1)} = B_k/p_k$. Coefficients A_k and B_k have been computed with data given in Part I and are plotted as functions of k . In the following the differential equation of heat transfer for an infinitely long fuel element ($\partial T^2 / \partial z^2 = 0$) is solved. Against the correct solution of the differential equation of temperature distribution in the fuel element $T(r,z)$ and of the longitudinal temperature distribution $t(z)$ in the coolant developed in Part I, axial heat transfer (in direction z) is neglected here. For this case, $\partial^2 T^* / \partial z^2 = 0$. This neglection considered, the corresponding approximate solutions $T^*(r,z)$ and $t^*(z)$ are obtained. Correct and approximate solutions are compared in the following. From graphs for the temperature distribution in three given cross sections of the fuel element and those for the longitudinal temperature distribution in the coolant, it is evident that differences between results of accurate and approximate calculations amount to a few % for the former, and only some hundredths of % for the latter. The error resulting from the neglect of axial heat transfer increases with the cross section of the fuel element.

Card 3/4

X

Steady-state heat distribution...

32721
II/008/62/000/001/002/002
B122/B102

and with its thermal conductivity, and decreases with its length. If the slenderness ratio of the fuel element $2R_0/l > 0.05$, and the thermal conductivity of its material $\lambda > 10 \text{ kcal/m}$, the more elaborate method is commendable, especially if the heat-flux density on the mantle is high. Differential equations have been solved for bare fuel elements only. There are 9 figures and 13 references: 12 Soviet-bloc and 1 non-Soviet-bloc.

ASSOCIATION: Központi Fizikai Kutató Intézet (Central Research Institute of Physics)

Card 4/4

X

SZABADOS, Laszlo, tudomanyos munkatars

Application of radioactive isotopes. Ipari energia 4 no. 6:
136-138 Je '63.

SZABADOS, Laszlo

Nuclear engineering news. Ipari energia 4 no. 7: 167-168
Jl '63.

1. "Ipari Energiagazdalkodas" szerkeszto bizottsagi tagja.

SZABADOS, Laszlo

Nuclear engineering news. Ipari energia 5 no.2:33 F '64.

1. Editorial board member, "Ipari Energiagazdalkodas."

SZABADOS, M.

Alga vegetation in the primordial moor at Kiskunhalas. p. 451, Vol 2, 1952. (published in 1954) A MAGYAR TUDOMAN EGYETEM BIOL GIAI INTEZETEINEK EVKONYVE. SZEGEDI RESZ. Budapest, Hungary.

So: Eastern European Accession. Vol 5, no. 4, April 1956

SZAFALCS, ...

SZAFALCS, ... Fauna and flora of some temporary stagnant pools in a meadow
of the Forzsony Mountains in Upper Hungary. II. Algae. In German.
p. 23.

Vol. 5, No. 3/4, 1954

WTA FICIGIA.

SCIENCE

Budapest, Hungary

So: East European Accession, Vol. 5, No. 5, May 1956

SZEGED, M.

"Life in the Tisza. II. Contributions to material on algae in the upper course of the Tisza." In German. p. 189.

ACTA UNIVERSITATIS SZEGEDIENSIS. PARS BIOLOGICA SCIENTIARUM NATURALIUM.
ACTA BIOLOGICA. Szeged, Hungary, Vol. 3, No. 3/4, 1957.

Monthly list of East European Accessions (EEAI), LC, Vol. 8, No. 8, August 1959.
Unclu.

SZABADOS, Pal, dr.

Development of planning methods for productivity in the construction industry. Munka szemle 6 no.10:13-18 0 '62.

SZABADOS, Pal, dr.

Manpower turnover in the construction industry and its effect
on the formation of productivity. Epites szemle 6 no.10:317-
322 '62.

1. Epitesgazdasagi es Szervezesi Intezet tudomanyos csoport-
vezetoje.

SZABADOS, Peter

Ferenc Rozsa, 1906-1942. Elet tud 16 no.5:131-134 29 Ja '61.

ABOSSY, Istvan, dr.; DOBOS, Laszlo, dr.; SZABADOS, Sandor, labor.
asszisztens

Clinical observations and early therapeutic results in
prednisolone treatment. Tuberkulozis 16 no.6:183-186 Je '63.

1. A satoraljaujhelyi Varosi Tbc Korhaz (igazgato: Abossy
Istvan dr.) kozlemenye.

(TUBERCULOSIS, PULMONARY) (PREDNISOLONE)
(CORTICOTROPIN) (SURGERY, OPERATIVE)
(ANTITUBERCULAR AGENTS)

L 10336-66 EWT(1)/EWA(1)/EWA(b)-2 RO/JK

ACC NR: AP6003344

SOURCE CODE: HU/0018/65/017/002/0153/0159

AUTHOR: Szabados, Terez-Sabadosh, T.; Boszormenvi, Jozsef-Besermen, I;
Dobias, Gyorgy-Dobiash, D.; Rojti, Mihaly-Ropty, M.; P. Juhasz, Vera-P. Yukhas,⁵³
ORG: HUMAN Institute of Vaccine Production and Research, Budapest (HUMAN)
Oltoanyagtermelo es Kutato Intezet); PHYLAXIA State Institute of Vaccine Production,
Budapest (PHYLAXIA Allami Oltoanyagtermelo Intezet)

TITLE: Changes in the antitoxin titer of animals used for serum production in the
course of iodocasein feeding

SOURCE: Kiserletes Orvostudomany, v. 17, no. 2, 1965, 153-159

TOPIC TAGS: experiment animal, biochemistry, veterinary medicine, animal physiology,
immunization, immunology

ABSTRACT: The changes in the antitoxin titer were studied in animals the metabolism
of which has been increased by iodocasein feeding. Among horses immunized with
diphtheria and tetanus toxin and sheep immunized with staphylotoxin, the antitoxin
level of the groups fed on iodocasein was significantly higher than that of the
controls. No noticeable difference was observed in cattle immunized with tetanus
toxin. Orig. art. has: 1 figure and 5 tables. [JPRS]

SUB CODE: 06 / SUBM DATE: 16Apr64 / ORIG REF: 003 / OTH REF: 002

Card 141

SZABC, L.; SZABADOS, Therese; ECK, Erna H., unter technischer Assistenz von
BERNATSKY, M.

Glutamic acid oxalacetic acid transaminase determinations in infancy
and childhood. I. Studies in relation to hydrocephalus. Acta Paediat
Acad Sci Hung 1 no.3:199-209 '60.

1. Kinderklinik der Medizinischen Universitat, Szeged.

(TRANSAMINASES blood) (HYDROCEPHALUS blood)

SZABO, L.; SZABADOS, Therese; ECK, Erna H., unter technischer Assistenz von
BERNATSKY, M.

Glutamic-oxalacetic acid transaminase determinations in infancy and
childhood. II. Studies on premature and newborn infants. Acta Paediat
Acad Sci Hung 1 no.3:211-221 '60.

1. Kinderklinik der Medizinischen Universitat, Szeged.

(TRANSAMINASES blood) (INFANT NEWBORN blood)
(INFANT PREMATURE blood)

SZABO, Lajos, dr.; SZABADOS, Terez, dr.; ECK, Erna, H. (technikai
munkatárs: BERNATSKY, Margit)

Glutamic acid-oxalacetic acid transaminase determinations in
infancy and childhood. Part II. Studies in connection with
hydrocephalus. Orv.hetil. 101 no.3:87-91 Ja '60.

(HYDROCEPHALUS diag.) (TRANSAMINASES chem.)

DUX, Erno, dr.; GIMESY, Ferenc, dr.; SZABADOS, Terez, dr.

Severe hemorrhagic diathesis after repeated exanguination-transfusion. Orv.hetil.101 no.33:1170-1174 14 Ag '60.

1. Szegedi Orvostudomanyi Egyetem, Gyermekklinika
(BLOOD GROUPS)
(BLOOD TRANSFUSION compl)
(HEMORRHAGIC DIATHESIS)

KOLTAY, Miklos, dr.; SZABADOS, Terez, dr.

Immunolectrophoresis. Orv.hetil. 101 no.37;1297-1303.11. S '60.

1. Szegedi Orvostudomanyi Egyetem, Gyermekklinika.
(ELECTROPHORESIS)

SZABADOS, Terez; BOSZORMENYI, J.; DOBIAS, G.; ROJTI, M.; JUHASZ, Vera P.

Effect of iodocasein feeding on the antitoxin titre of animals used in serum production. Acta microbiol. acad. sci. Hung. 10 no.4:387-396 '63-'64

1. Serum and Vaccine Institute "Human" (Director: G. Veres), Budapest, and Serum and Vaccine Institute "Phylaxia" (Director J. Molnar), Budapest.

SZABADOS MOLNAR, Arpad

Useful technique. Radioteknika 11 no.11:345 N '61.

STANKOVIANSKY, Samo, prof., inz.; RUSINA, Rudolf, inz., C.Sc.; SZABADOSOVA,
Katarina, promovana chemicka

Polarographic examination of isothiocyanates nonsoluble in water
(I). α -naphthylisothiocyanates and β -naphthylisothiocyanates.
Chem. zvesti 16 no.8:625-632 Ag '62.

1. Katedra analytickej chemie, Prirodovedecka fakulta University
Komenskeho, Bratislava, Smeralova 2.

RUSINA, R. [deceased]; STANKOVIANSKY, Samo, prof., inz.; SZABADOSOVA,
Katarina, promovana chemicka

Polarographic examination of isothiocyanates nonsoluble in
water. Pt. 2. Chem zvesti 17 no.5:300-306 '63.

1. Katedra analytickej chemie, Prirodovedecka fakulta University
Komenskeho, Bratislava, Smeralova 2.

SABADVARI, Ferents [Szabadvary, Ferenc]

History of sugar. Nauka i zhizn' 29 no.10:90-91
0 '62. (MIRA 15:12)
(Sugar)

"APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001654320017-2

SZABADVARY, Ferenc

History of aluminum. Elet tud 16 no.29:899-902 16 Jl '61.

APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001654320017-2"

"APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001654320017-2

SZABADVARY, Ferenc

Darivatograph: a new Hungarian invention. Elet tud 18 no.29:
920-922 21 Jl '63.

APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001654320017-2"

SZABADVARY, F.

HUNG.

✓ Ascorbometric determination of oxygen dissolved in water. L. Erdey and F. Szabadvary (Tech. Univ., Budapest). *Acta Chim. Acad. Sci. Hung.*, 4, 325-32 (1954) (in German) (English summary); cf. preceding abstr.—To det. dissolved O₂ in H₂O, treat the sample with NaOH soln. and a soln. of Fe(NH₄)₂(SO₄)₂ that has previously been passed through a Cd reductor to remove traces of Fe⁺⁺⁺. Dissolve the ppt., contg. Fe⁺⁺⁺ in acid, and titrate the Fe⁺⁺⁺ with 0.01*N* ascorbic acid, with 4-amino-4'-methoxydiphenylamine indicator. NO₂⁻ interferes with the detn.

A. L. Underwood

Szabóvary F.

8. Determination of oxygen dissolved in water by means of ascorbic acid - *Vizben oldott szerves mérőszállító aszkorbinsavval* - I. Boddy and E. Szabóvary - Journal of Hydrology - *Hydrologie* - Vol. 36, 1954, No. 1-2, pp. 8-14, 2 figs., 2 tables.

Oxygen dissolved in water was determined by filling a flask of known volume with the water to be tested. By means of a long pipette thin layers of an iron(III) free iron(II) ammonium sulfate solution and a sodium hydroxide solution were measured at the bottom of the flask. The iron(II) solution was made iron(III) free by passing it through a Cd-reductar pipette or by shaking with iron powder. Iron(III) formed by the action of the dissolved oxygen -- after shaking, settling and acidification -- was titrated with an 0.01 N ascorbic acid measuring solution in the presence of 4-amino-4'-methoxy-diphenylamine as an indicator. Winkler's method of oxygen determination may also be modified by adding iron(II) ammonium sulfate instead of potassium iodide to the oxidized manganese solution, and by titration with ascorbic acid.

M

SZABADVÁRY, F.

4-amino-4'-methoxy-diphenylamine as a colorimetric reagent. I. Determination of iron(III) ions — I.
Pádey, F. Szabadváry. *Magyar Kémiai Folyóirat*, Vol. 62, No. 3, pp. 311-316, 4 figs., 9 tables.)

The 4-amino-4'-methoxy-diphenylamine compound is transformed into a blue-coloured product in the presence of oxidizing ions. This colour reaction was employed for the determination of iron(III) ions. It was found that an inexpensive textile dyestuff, Virtämin blue B, produced on an industrial scale was suitable as a reagent. The method proved to be useful for the determination of 0.2 to 12.0 µg/ml iron(II) in the pH range of 1 to 4. In the presence of other oxidizing ions yielding a blue colour reaction with the reagent, first the total extinction of the solution was measured and subsequently the measurement was repeated after the addition of sodium citrate. The difference between the two values obtained corresponded to the extinction of the iron(III) ions.

SZABADVARY, FERENC

5
9
0

✓ 4-Amino-4'-methoxydiphenylamine as a colorimetric reagent. Determination of iron(III) ions. László Bardey and Lajos Szabadvary. Tech Univ. Budapest. Acta Chim. Acad. Sci. Hung. 6, 131-142 (1963) in German with English summary; cf. C.A. 47, 8295t. — The blue color, with absorption max. at 570 m μ , formed in the presence of oxidizing agents by 4-amino-4'-methoxydiphenylamine, available as Vaseline Blue B Base, is suitable for the detn. of 0.02-12 p.p.m. of Fe(III) in the pH range 1-4 at temps. from 10 to 30°. Ions such as citrate, tartrate, $\text{P}_2\text{O}_7^{4-}$, CaO_4^{4-} , F^- , and PO_4^{3-} interfere b' preventing color formation. There is no interference by Na^+ , K^+ , NH_4^+ , Mg^{2+} , Ca^{2+} , Ba^{2+} , Mn^{2+} , Ni^{2+} , Co^{2+} , Al^{3+} , Cu^{2+} , As^{3+} , Sb^{3+} , Cl^- , Br^- , I^- , SO_4^{2-} , NO_3^- , or AsO_4^{3-} . Bi^{3+} , Se^{4+} , Fe(CN)_6^{4-} , and Fe(CN)_6^{3-} must be absent, whereas certain amts. of Fe^{2+} , Zn^{2+} , Cd^{2+} , Pb^{2+} , CN^- , and BO_4^{3-} are permissible. In the presence of oxidizing agents the total oxidizing power is detd. and then the Fe is complexed with citrate and the amt. of Fe(III) detd. by difference. B. P. Block

Szabadvary, F.

Colorimetric determination of iodine with the use of
Verlamine Blue. L. Erdey and F. Szabadváry (Tech.
Univ., Budapest). *Acta Chim. Acad. Sci. Hung.* 1961, 14, 101.

Univ., Budapest). *Zs. Zool.* Acad. Sci. Hung. 8, 191-202 (1955) [in German] (English summary).—Attempts to develop a colorimetric I detn. (e.g. the reaction of I with starch) were not successful. However the blue color of I with Vanlamine Blue (4-amino-4'-methoxydiphenylamine) could be developed into a colorimetric I detn. of great sensitivity. Procedure: Transfer the weakly acid soin. contg. 23-000 γ of I dissolved in not more than 25 ml. H₂O to a 50-ml. volumetric flask. Add an acetic acid-acetate buffer soln. of pH 3 and finally 2 ml. of a 1% soln. of Vanlamine Blue. After waiting 2 min., measure the color with a Pulfrich photometer and filter S 57 at water. A pH of 3 favors act only greater color stability but prevents interference by iodide. The absorption curve follows Beer's law quite closely over a pH range of 0-5, except for I concns. under 12 γ /ml. Concns. of 0.6 γ /ml. I could be accurately detd. by this colorimetric method. Oxidizing as well as reducing substances must be absent, also Ph, Ig, Bi, CN, oxalate, and borate. 14 references. Ernst M. Goldstein

Ernest M. Goldstein

APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001654320017-2"

Szabadváry, Ferenc

W

74285* Variumin Blue B as a Colorimetric Reagent. Vari-
aminkék (4-amino-4'-metoxi difenilamin) mint kolori-
metriás reagens. II. Determination of Iodine. Jódmeg-
határozás. (Hungarian.) László Erdéy and Ferenc Szabadváry.
Magyar kémiai folyóirat, v. 81, no. 11, Nov. 1935, p. 841-845.

①

A color reaction suitable for the determination of iodine in
solution of 0.5-13 µg/I per ml. between the pH range of 1-5.
Sensitivity of method. Tables, graphs. 15 ref.

AB 500

SZABADVARY, F.

-7

6/25/83

VAN DER HORST, B. A. ~~in communication~~
Analyst, Bureau of Mineral Resources
Geological Survey of Canada

July 1957, 68 (6), 153-158
blue B as an indicator, in the pH range 3 to 4
0.5 to 12 µg per ml of V^{IV} and 0.2 to 4 µg per ml of
Cr (as chromate) can be determined photometrically.
The oxidation of V and Cr to the required
valency states is described. Iron, Cr and V can be
determined simultaneously by the same method.
The theory of these determinations and the effect
of interfering ions are discussed. A G Pinto

PD 11/13

SZABADYARY, F.

Cham 3476. *Varianin blue as a colorimetric reagent.* L. Erdéy and F. Seabadvity (Inst. für Allgemeine Chemie, Tech. Univ., Budapest). *Z. anal. Chem.*, 1937, 155 (2) 90-96.—Varianin blue can be used for the colorimetric determination of many ions whose standard potential lies above that of the reagent, e.g. H^+ , Cl^- , Mn^{2+} , V^{2+} , Ag^{+} , I^- , and Fe^{2+} . Oxidation equivalents of different ions give the same extinction reading, so that one calibration curve suffices. An aq. soln. of the reagent (0.5%) (2 to 6 ml.) is added to the test soln. 0.5 to 20 μg equiv. in water or alcohol. The extinction is measured after 5 min. with a Pulfrich 537 filter (370 m μ). The mechanism of formation of the blue compound, the effect of changes of conditions, and interferences are discussed. A procedure is described for the estimation of H^+ , Mn^{2+} , V^{2+} , and Fe^{2+} in the presence of each other. G. B. Wicks

APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001654320017-2"

"APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001654320017-2

APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001654320017-2"

Szabadvári

Distr: 4E2c

24. A photometric method of determining vanadium and chromium with Variamine Blue. (In German), I. Bidey, F. Szabadvári, Acta Chimica Academiae Scientiarum Hungaricae, Vol. 13, 1958, No. 3-4, pp. 335-345, 2 figs., 7 tabs.

Though several methods are known for the colorimetric determination of vanadium, they have the common fault that the specific extinction is small. Considerably better results can be obtained with Variamine Blue. Vanadium(V) may be reliably determined within the concentration range of 0.5-12 $\mu\text{g}/\text{ml}$ whereas chromium in that of 0.2-4 $\mu\text{g}/\text{ml}$ in the presence of each other in solutions of pH 1-4. Potassium permanganate is used for oxidizing vanadium and potassium persulfate for chromium, the excess of these reagents is removed before adding the Variamine Blue. In addition a simple colorimetric procedure is given for the determination of iron, vanadium and chromium in the presence of one another by means of Variamine Blue.

LB

SZABADVARY, FERENC

5
1-89 (WA)

Oxidation products of 4-amino-4'-methoxydiphenylamine.
Éva Bányai, László Erdéy, and Ferenc Szabadváry (Tech. Univ., Budapest). *Acta Chim. Acad. Sci. Hung.* 20, 307-20 (1959) (in German).—The polarographic waves and the absorption max. of 4-amino-4'-methoxydiphenylamine (I), of its oxidn. products, and of *N*-(*p*-anisyl)-*p*-benzoquinone d-imine (II) proved that II formed in the 2-electronic oxidn. of I. By varying the pH value of the soln., II suffered a change of color, due to the proton affinity of the imino group. The degree of proton affinity of the imino group was established by an optical method and on the basis of the break points of the oxidn.-redn. potential: pH curves. The electrode potential of the oxidn.-redn. system proved to be pH dependent. In a slightly acidic medium, oxidn. took place through a semiquinone intermediate (III), as detd. by using the index potentials. In the oxidn.-redn. potential measurements, the oxidn. agents were: 0.01*N* Br-H₂O (in acidic soln.) or 0.01*N* K ferricyanide (in alk. soln.), resp. During the potentiometric oxidn. of I with Br-H₂O at pH 1-6, I gave at first a blue color. By adding Br-H₂O in an amt. corresponding to 2 electrons a violet color arose; and in the presence of strong oxidizing agents (Br-H₂O and Cl-H₂O in great excess), the soln. became red. Over pH 8 the oxidized soln. was continuously yellow.

At pH 1.5-5.5, the 2-electronic oxidn. went through the intermediate III, the stability of which was assured by mesomeric structures. In alk. soln. the oxidn. was direct. At pH 3, a protonated form of II (IV) presented an absorption max. at 580 m μ . The pH region 3-4 was the most favorable for IV (25%). The 2-electronic oxidn. product of I was violet in acidic soln. (absorption max. at 540 m μ), red close to pH 7 (max. at 460 and 540 m μ), and yellow in alk. soln. (max. at 460 m μ); consequently the red color was a mixed one. By polarographic and optical methods, this oxidn. product proved to be II. The color change was explained as follows: in alk. soln., II exists as a yellow base; in acidic soln., however, by taking up a proton, II can exist in the two violet mesomeric forms of IV. The overoxidized product arising from the action of Cl-H₂O contained 3.1% N, no Br; and, probably, it was decompd. Below pH 1, the violet IV became colorless by decompn. into *N*-(*p*-anisyl)-*p*-benzoquinone imine and NH₃. In weakly acidic medium I took up only one proton, probably on the primary amino group. Over the pH range 1-6, therefore, both the oxidized and the reduced forms of I may exist as univalent cations.

E. Kasztreiner

SZABADVARY, Ferenc (Budapest)

"Determination of phlogiston dissolved in water." Kem tud kozl MTA
16 no.1:41-46 '61.

1. Budapesti Muszaki Egyetem, Altalanos Kemialos Tanszek.

(Chemistry—Phlogiston)

SZABADVARY, Ferenc

The first chemical analysis of the Buda mineral waters.
Magy kem lap 17 no.5:217-220 My '62.

1. Budapesti Muszaki Egyetem Altalanos Kemialos Tanszeke.

CHARLOT, G.; SZABADVARY, Ferenc[translator]

Some examples for the role of reactions occurring in a solution in
the further development of chemical analysis. Kem tud kozl MTA 16 no.3:
247-254 '61.

DUVAL, Cl.; SZABADVARY, Ferenc[translator]

Thermogravimetric and infrared spectrophotometric investigation of solid-phase reactions. Kem tud kozl MTA 16 no.3:255-264 '61.

1. Ecole Nationale Superieure de Chimie, Paris.

SZABADVARY, Ferenc

Beginnings of chemical industries. Elet tud 16 m.16:483-486 16
Ap '61.

SZABADVARY, Ferenc

What is behind the tabulation of Mendeleev. Elet tud 16
no.53:1685-1687 31 D '61.

BECK, Mihaly; BITE, Pal; BRUCKNER, Gyozo; CSENTES, Jozsef; CSUROS, Zoltan;
DEAK, Gyula; ERDEY-GRUZ, Tibor; ERDEY, Laszlo; FABIAN, Pal;
FINALY, Istvan; FODOR, Gabor; FODORNE CSANYI, Piroska;
GYORBIRO, Karoly; INZELT, Istvan; KUCSMAN Arpad; NEUMANN, Erno;
PUNGOR, Erno; SCHNEER, Anna; SCHULEK, Elemer; SZABADVARY, Ferenc

Rules for the Hungarian chemical nomenclature and orthography.
Kem tud kozl MTA 17 no.1/4:1-292 '62.

1. "A Magyar Tudomanyos Akademia Kemial Tudomanyok Osztalyanak Kozlemenyei" szerkeszto bizottsagi tagja (for Bruckner, Csuros, Laszlo Erdey, G.Fodor, and Schulek). 2. "A Magyar Tudomanyos Akademia Kemial Tudomanyok Osztalyanak Kozlemenyei" szerkesztoje (for Erdely-Gruz). 3. "A Magyar Tudomanyos Akademia Kemial Tudomanyok Osztalyanak Kozlemenyei" technikai szerkesztoje (for Finaly). 4. Muvelodesugyi Miniszterium (for Csentes). 5. Magyar Tudomanyos Akademia Helyesitasi Bizottsage (for Fabian). 6. Nehezipari Miniszterium (for Neumann).

HAISSINSKY, M.; SZABADVARY, Ferenc [translator]

Comparison of the chemistry of cis and trans-uranides with that
of lanthanides and of transition elements. Kem tud kozl 18
no.3:395-411 '62.

1. Faculte de Sciences, Laboratoire Curie, Paris (for Haissinsky).

SZABADVARY, F.

"Lexicon of chemical-pharmaceutical symbols" by Wolfgang
Schneider. Acta chimica Hung 32 no.1:129 '62.

SZABADVARY, F. (Budapest, XI., Gellert ter 4)

Founding of the Selmec (Schemnitz) Mining Academy and its
significance in teaching chemistry all over the world. Periodica
polytechn chem 7 no.2:127-133 '63.

1. Lehrstuhl fur Allgemeine Chemie, Technische Universitat,
Budapest.

SZABADVARY, Ferenc

Paul Szily a forgotten Hungarian scientist. Orv.hetil. 105 no.7:
328-329 16 F '64.

1. Budapesti Műszaki Egyetem, Altalanos Kemiai Tanszek.

SZENTIVANYI, J., KOMORNÍK, TIBOR, A.

Relationship between quick variations in the geomagnetic field and telluric currents. p. 544. (SÁVYASZATI LAPOK, Budapest, Hungary), Vol. 9, No. 10, Oct. 1954.

SC: Monthly List of East European Accessions, (EEAI), 1C, Vol. 4, No. 5, May 1955. Uncl.

SZABADVARY, Laszlo; SZABO, Margit

Geoelectric bauxite prospecting measurements in the Bakony
Mountains. Geofiz kozl 13 no.3:263-272 '64.

SZABADVARY, Laszlo

Geoelectric soil research on the basis of test borings. Geofiz.kozl
3 no.1/11:121-141 '54.

SZABADVARY-L.

EV
②
3
57. Correlations between the rapid variations in the geomagnetic field and the telluric currents - E. Anna and A. Erkel. L. Szabadváry. (Műszaki Lapok - Vol. 9 (87), 1956, No. 10, pp. 544-549, 11 figs.)

The authors have designed a magnetometer with a sensibility of $10^{-8} \gamma$, with which rapid magnetic variations have been measured, and simultaneously the variations of the telluric currents have also been recorded. The measuring circuit is a circular conductor of 400 m dia, consisting of two turns, the dimensions of which can be increased up to a certain limit for increasing the sensibility of the apparatus. The recording device records the N and E components of the telluric currents as well as the voltage induced in the circular conductor on a photoreel. The two components are recorded by two separate galvanometers, a third galvanometer records the induced voltage. For the determination of the voltage variations recorded on the photoreel a calibrating device is required by which the two poles of the galvanometer receive a voltage of 7.82×10^{-4} volt from a 1.5-volt battery through a double potentiometric connection. It is indispensable that the measurements be executed in an undisturbed area; this can be found in the mountains surrounding Sopron. The results of the measurements show that between the rapid variations of the magnetic field and the telluric currents there exists a certain relationship which can be utilized in applied geophysics. The higher the voltage induced by telluric currents in a circular conductor arranged in a horizontal plane, the smaller

E. ANNUAU

the angle formed by the direction of the inclination of the base rock and the horizontal projection of the direction of inclination of the telluric currents, and the greater the angle of inclination of the base rock. The reverse is true for a vertical circular conductor. For the combined application of both circular conductors, a procedure indicating the depth, the strike line and the angle of inclination of the base rock can be elaborated. On the basis of these relations it becomes possible to employ several methods of geophysical research.

3

7/2

SZABADVARY, Laszlo

Geoelectrical exploration experiences with the location of water wells in Mongolia. Geofiz kozl 9 no.3/4:135-164 '61.

(Mongolia--Water) (Geology)

HOBOT, Jozsef; BRKEL, Andras; SZABADVARY, Laszlo

Complex geoelectric measurements for basin exploration in South
Transdanubia. Geofiz kozl 13 no.3:273-288 '64.

"APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001654320017-2

ERKEL, Andras; KIRALY, Erno; SZABADVARY, Laszlo

Instrument series of GE type geoelectric resistance meters. Geofiz
kozl 13 no.1:71-82 '64.

APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001654320017-2"

SZABADY, Egon (Budapest)

Tasks of the Demographic Committee, Hungarian Academy of Sciences.
Magy tud 68 no.3:175-180 Mr '61. (EEAI 10:6)

1. Foosztalyvezeto, Kozponti Statisztikai Hivatal, Budapest.
(Hungarian Academy of Sciences) (Demography)

SZABADY, Egon, Dr.

The session of the International Union for the Scientific Study of
Populations. Stat szemle 40 no.1:79-80 Ja '62.

SZABADY, Egon

Demographic Symposium. Magy tud 70 no.2:121-123 F '63.

1. Kozponti Statisztikai Hivatal foosztalyvezetoje.

SZABADY, Egon

Delegation of the Hungarian Research Group on Demography
in Stockholm. Magy tud 70 no.10:716-718 O '63.

1. Kozponti Statisztikai Hivatal foosztalyvezetoje.

CSIKOS, Bela; FUTO, Istvan; EROS, Jozsef; SZABADY, Jeno; EISLER, Janos, Dr.;
WALLENSTEIN, Mihaly; REMBECZKY, Laszlo; BALINT, Gabor;
ASZTALOS, Peter; BERENyi, Laszlo, okl. gepezmernok;
HORCHER, Frigyes

Remarks on the article "The most important problems of
technical development and network electrical installations
and tasks for the manufacturing industry related to this."
Villamossag 9 no.1/3:17-23 Ja-Mr '61.

1. Az Eromu Troszt villamos osztalyanak vezetoke (for Csikos).
2. A Nehezipari Miniszterium Villamosenergiaipari Igaggatisaganak Szakosztalyvezetoe (for Futo). 3. VERTESZ Villamos Eromu Tervezo es Szerelo Vallalat (for Eros). 4. Klement Gottwald Villamossagi Gyar (for Szabady, Wallenstein, Rembeczky, Balint, Asztalos, Horcher). 5. Budapesti Muszaki Egyetem (for Eisler).

FARKAS, Bela; HOLCZHAUSER, Albert; FUREDI, Pal; SZEPESI, Endre, Dr.;
SZABADY, Jeno; SZEPESSY, Sandor; HALASZ, Antal; BALLAI, Laszlo;
SZEKELY, Istvan; KOHUT, Matyas

Remarks on the article "Problems of technical development for the heavy industry on the basis of the requirements of industrial branches which use its products." Villamossag 9 no.1/3:53-61 Ja-Mr '61.

1. A Klement Gottwald Villamossagi Gyar fomernooke (for Farkas).
2. A TRANSZVILL Transzformator es Villamoskeszulekgyar fomernooke (for Holczhauser).
3. VERTESZ Villamoseromu Tervezo es Szerelo Vallalat (for Furedi).
4. Hoenergiagazdasagi es Tervezo Vallalat (for Szepesi).
5. Klement Gottwald Villamossagi Gyar (for Szabady and Szekely).
6. Csepeli Transzformatorgyar (for Halasz). 7. Ganz Kapcsolok es Keszulekek Gyara (for Kohut).

SZABADY, Jeno, okleveles gepeszernok

Modern heavy-current insulators. Technika 6 no.3:6-7 Mr '62.

SZABADY, Karoly

Tasks of the mechanization of the construction industry. Ujít lap
14 no.19:10-11 10 0 '62.

SZAFAN, J.

"Tasks of Light Industry in Standardization," P. 297. (WIADOMOSCI, Vol. 22,
No. 6, June 1954. Warszawa, Poland)

SO; Monthly List of East European Accession, (EEAL), LC, Vol. 4,
No. 1, Jan. 1955 Uncl.

Castany, J.

Questions of insulation of electric machines. p. 3., (VILMOSZKI, Budapest, Hungary), Vol. 2, No. 1/2, Jan./Feb. 1954.

EE: Monthly List of East European Accessions, (EAA), 1G, Vol. 4, No. 5, May 1955.

SZABADY, Jeno

The Klement Gottwald Electrical Works are 80 years old.
Elektrotechnika 52 no.3:127-133 '59.

1. A Klement Gottwald Villamossagi Gyár fomernöke.

SZARLA, Jan

The RHL3 power loader. Przegl kolej mechan 15 no.1:30-31 Ja '63.

1. Zaklady Naprawy Taboru Kolejowego, Wroclaw.

SZABE, Istvan; SOMFAI, Karoly

Up-to-date methods for treating hardened materials.
Ujít lap 14 no.9:58-60 10 My '62.

SZABLEWSKA, Anna

Difficulties in the adaptation of children in an anti-tuberculosis sanatorium and their causes. Pediat. pol. 38 no.10:889-895 O '63.

1. Z Dziecięcego Ośrodka Sanatoryjno-Prewentoryjnego w Rabce
Dyrektor: dr med. J. Rudnik.
(HOSPITALIZATION) (TUBERCULOSIS IN CHILDHOOD)
(ADAPTATION, PSYCHOLOGICAL)

BASINSKI, Antoni, SZYMANSKI, Wojciech, SZABLEWSKI, Lech

Solubility of thallium ferrocyanide in water and in some organic solvents as determined by the tracer method. Roczniki chemii 36 no.7/8:1255-1257 '62.

1. Katedra Chemii Fizycznej, Uniwersytet im. M.Kopernika,
Toruń.

SZABLEWSKI

✓ 1953. Stablewski, E., Turbulent flow in divergent channels (in German), Ing.-Arch. 22, 4, 268-281, 1954.

Using Prandtl mixing-length assumption to relate turbulent shear stress to mean velocity, author analyzes flow in a two-dimensional channel of small divergence (less than 5°). As a first approximation, the velocity gradient terms in the boundary-layer equations are neglected and velocity distributions are derived for regions close to and far from the wall. These are variations of the well-known logarithmic and wall laws. Comparison is made of the derived laws to the results of Kurnazarov (VNI / Zeschriften 80, 1926). Trend is very x in agreement, but numerical results are not. A second approximation is obtained by combining the mixing-length assumption with a modified momentum equation. The resulting integral equation is linearized to obtain an explicit solution. Good agreement is found between the second approximation and experimental results.

In spite of verification of the theoretical laws by the experiments, reviewer does not think paper makes a substantial contribution to boundary-layer theory. Mixing-length distribution is arbitrary and has no fundamental importance. Recent advances in the analysis of boundary layers under pressure gradient should eliminate the need for analyses such as are presented in this paper. E. D. Eaines, Canada

P. E. E.

SZ. A. GLEWSKI, W.

2

2608. Szyldowicz, W., Velocity distribution close to the wall
in boundary-layer flow with pressure increase (in German), Ing.-
Arch. 23, 4, 295-306, 1953.

AL Phip
Lilley

The mean equations of motion for turbulent flow near a wall in
the presence of a pressure gradient have been integrated to find
the mean velocity distribution (law of the wall) using the Prandtl
mixing-length approximation for the turbulent shear stress. The
constants in the mean velocity formula are assumed to be equal
to universal constants given by the empirical constants in the
zero pressure gradient case. Comparison with results of
Nikuradse's experiment for plane diffusers shows fair agreement,
including the diffused angle for separation, which was found by
extrapolating the wall shear stress to zero. Comparison with more
recent experimental results is not given.

G. M. Lilley, England

R.H.A.
MT